URBAN HEAT ISLAND ANALYSIS

SEATTLE, WASHINGTON | 2020

EARTH ECONOMICS

THE DANGER OF EXTREME HEAT

Extreme heat – exacerbated by the "urban heat island" effect and climate change - is the deadliest weather or climate related hazard in the U.S. Urban heat islands occur in areas with a high percentage of impervious surface area and little green space where temperatures can significantly exceed the regional average. These heat islands are found in dense urban areas with a limited greenspace and large amounts of asphalt, concrete, and other impervious surfaces.



Global temperatures have risen approximately 1°C over the past century.

|--|

Urban areas have risen an additional 1-3°C due to the urban heat island effect.


ANALYSIS FOR SEATTLE, WASHINGTON

In Seattle, our analysis found that urban heat islands increase maximum temperatures by as much as **6° F** above the local baseline. Combined with regional climatic effects, neighborhoods in Central and South Seattle may be as much as 14° F hotter than neighborhoods in North Seattle (Map 1). **33,000** people live in neighborhoods where maximum temperature exceed the 80th percentile and where average per capita incomes are less than 200% the federal poverty level (identified by black outline in the map below). **10,300** people live in areas that meet the above criteria, have a high percentage of population in high risk age groups (under age 14 or over age 65), and exceed 50% of regional population density. Every additional degree carry significant public health costs.²

In the Seattle's most severe urban heat islands, all-cause mortality rates are estimated to increase by **as much as 7.7%** when regional temperatures exceed 82°F. Seattle experiences 25 days above 82° per year, on average. Our model estimates that **8 – 15 lives** are lost in the city each year due to extreme heat exacerbated by urban heat islands.

The public health burden of urban heat islands is disproportionately levied on the lowest income households. In Seattle there is a strong correlation between household income and urban heat islands (Graph 2).

Without intervention, Seattle's heat island impacts will grow significantly in a changing climate. Number of days above the 82° F threshold has increased by more than 50% since 1980 (Graph 3). On the current trajectory Seattle will experience 42 days above the temperature threshold by 2050, and 59 days above the threshold by 2100.



WITH A NATURE-BASED SOLUTION

Urban heat islands are a critical public health issue – with a nature based solution. Urban trees and greenspace can effectively mitigate urban heat islands, while providing numerous benefits such as air quality improvement and stormwater capture.³



Urban vegetation offers shade and evapotranspiration, which can lower peak summer

temperatures 2-9°F.^{3,4}



Urban trees provide heat reduction, which supports public health by reducing heat stress and contributes to reduced household energy costs. THERE IS A STRONG **CORRELATION BETWEEN** SUMMER HEAT AND NEGATIVE HEALTH OUTCOMES.



TEMPERATURES ABOVE 82°F SIGNIFICANTLY **INCREASE THE RISK OF** CARDIOVASCULAR DISEASES. **RESPIRATORY ILLNESSES,** AND HEAT STROKE.

WHAT ARE **URBAN HEAT ISLANDS?**


Dense urban areas absorb and trap heat due to impervious surfaces — like pavement and rooftops that absorb solar radiation, buildings that block wind flow, pollution and smog that trap solar radiation, and low levels of vegetation and tree cover that restrict evapotransporation — which functions as natural air conditioning.

Urban heat islands have significant impacts on human health and will have an even greater impact over the next century as the climate warms and populations urbanize.


EARTH ECONOMICS TAKING NATURE INTO ACCOUNT

107 N. Tacoma Avenue Tacoma, WA 9840 eartheconomics.org

THE COST OF **PUBLIC HEALTH CAUSES THE GREATEST ECONOMIC BURDEN ON LOW-INCOME COMMUNITIES.**



TEMPERATURES IN SEATTLE **HAVE INCREASED BY** MORE THAN 50% IN THE LAST 20 YEARS.



REFERENCES

- Anderson, B. G., & Bell, M. L. (2009). Weather-related mortality: how heat, cold, and heat waves affect mortality in the United States. Epidemiology (Cambridge, Mass.), 20(2), 205.
- ² Department of Environmental Health, Harvard T.H. Chan School of Public Health. 2016. Study on the Association Between Ambient Temperature and Mortality Using Spatially Resolved Exposure Data. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5071163/
- ³ Wang, Y., Akbari, H., 2016. The Effects of Street Tree Planting on Urban Heat Island Mitigation in Montreal. https://doi.org/10.1016/j.scs.2016.04.013.

